A Molecular Orbital Study on the Interaction Between BF₃ and the Compounds Dimethyl Ether and Perfluorodimethyl Ether

Wilfredo Morales Lewis Research Center Cleveland, Ohio

March 1996



A MOLECULAR ORBITAL STUDY ON THE INTERACTION BETWEEN BF₃ AND THE COMPOUNDS DIMETHYL ETHER AND PERFLUORODIMETHYL ETHER

Wilfredo Morales
National Aeronautics and Space Adminstration
Lewis Research Center
Cleveland, OH 44135

ABSTRACT

A molecular orbital method was used to study the interaction between BF_3 (a Lewis acid) and the compounds dimethyl ether and perfluorodimethyl ether. The results indicated the existence of a Lewis acid-base interaction between the BF_3 molecule and dimethyl ether but no interaction with the perfluorodimethyl ether. In addition, the results of a second molecular orbital study, involving the interaction between AlF_3 (another Lewis acid) and perfluorodimethyl ether, also indicated no Lewis acid-base interaction.

INTRODUCTION

Many researchers (refs. 1 to 4) have investigated the high temperature catalytic decomposition of perfluoropoly-alkylethers (PFPEs) in the presence of metal oxides and metal fluorides in order to explain, and hopefully prevent, PFPE degradation under tribological conditions. Two viewpoints have emerged concerning the decomposition mechanism. Some researchers (refs. 1, 2, and 5) have postulated that the surface Lewis acid sites of certain metal oxides and metal fluorides attacked the oxygen lone-pair electrons of the PFPEs initiating decomposition. Other researchers (refs. 3, 6, and 7), however, have conducted experiments suggesting that these surface Lewis acid sites may not be involved in the decomposition of PFPEs.

Ball and Zehe (ref. 8) added support to the latter viewpoint by conducting infrared matrix-isolation spectroscopy studies on the interaction between BF_3 (a strong Lewis acid) and the model compounds $(C_2H_5)_2O$ and $(C_2F_5)_2O$. Their findings confirmed the known Lewis acid-base interaction between BF_3 and the diethyl ether but no observable interaction with the perfluorodiethyl ether. The possibility, however, was raised that infrared-active vibrations due to BF_3 -perfluorodiethyl ether interaction may have been too weak for detection under the co-condensed conditions of the matrix isolation apparatus.

A computational molecular orbital (MO) study on the interaction between BF₃ and two model compounds, dimethyl ether (DME) and perfluorodimethyl ether (PFDE), was undertaken to hopefully reproduce (computationally) the experimental results of Ball and Zehe, using two similar but smaller model ether molecules. The interaction between AlF₃, another Lewis acid, and PFDE was also studied since the catalytic decomposition of PFPEs by AlF₃ at elevated temperatures is known to occur (ref. 1).

MODEL COMPOUNDS AND COMPUTATIONAL DETAILS

The AM1 self-consistent field molecular orbital procedure developed by Dewar and co-workers was used in this study. Although <u>ab initio</u> methods can be used for these small molecules, it is our intent to study the behavior of very long chain perfluoro ether molecules in the near future. The use of <u>ab initio</u> methods would become prohibitively time consuming and calculationally expensive for these exceedingly large molecules. It was thus decided to employ semiempirical methods (AM1 or PM1) for current and future work.

The computations employed in this study used the HyperChem version 4.0 of AM1 using the RHF option on a 486DX personnel computer. Complete geometric optimization during energy minimization was performed for all single molecules and for all interacting pairs of molecules. Configuration interaction was not used. Computing times ranged from 1 to about 8 hr.

RESULTS AND DISCUSSION

Before computational runs were performed on the interaction between the ether molecules and the Lewis acid molecules, two preliminary tests of the AM1 program were made. The first test compared the calculated proton affinity of the perfluorodimethyl ether molecule, using the AM1 program, with the proton affinity calculated using ab initio methods. The second test involved the analysis of the computational results between the interaction of NH₃ and BF₃.

First Test

An initial molecular geometric structure for PFDE was obtained using the model builder routine of the semi-empirical program (fig. 1(a)). Figure 1(b) shows the optimized molecular structure and atomic charges of PFDE as predicted by the AM1 method. Table I lists the calculated bond lengths, angles, and dihedral (or torsion) angles of this molecule and compares them to the calculated values obtained by other researchers using RHF/3-21G (ref. 9) and RHF/4-31G (ref. 10) ab initio methods. The experimental values (ref. 10) are also listed. The bond lengths and angles calculated using the AM1 method are less than 3 percent greater than the experimental values. Both the AM1 and ab initio results are in excellent agreement with the experimental results. The dihedral angle predicted by AM1 was 1.4° greater than the experimental value of 164°.

Vibrational analysis of the AM1 optimized structure revealed no negative frequencies indicating this was indeed a valid minimun energy structure.

The optimized PFDE molecule was then protonated by interacting H^+ with the O atom of the ether molecule. Figure 2 shows the optimized protonated ether molecular structure. Vibrational analysis revealed no negative frequencies. Table II lists and compares the bond lengths and angles with the values obtained by the RHF/3-21G <u>ab initio</u> method (ref. 9). Proton affinity (PA) is defined as the energy given off when a molecule accepts a proton in the gas phase:

$$M + H^{+} \rightarrow MH^{+}$$
 $\Delta H \equiv \text{proton affinity}$ (1)

The proton affinity can thus be calculated using heats of formation in the following equation:

$$\Delta H_{rxn} = \Delta H_f(MH^+) - \Delta H_f(M) - \Delta H_f(H^+)$$
 (2)

where:

Proton Affinity (PA) = $-\Delta H_{rxn}$ ΔH_f of Protonated Perfluorodimethyl ether (MH⁺) = -671.8 kJ/mol (this work) ΔH_f of Perfluorodimethyl ether (M) = -1592.7 kJ/mol (this work) ΔH_f of H⁺ = 1537.4 kJ/mol (experimental value)

The PA value calculated using equation 1 was 616.4 kJ/mol compared to the PA value of 585.5 kJ/mol calculated by Fontaine et al. (ref. 9). The AM1 PA result was thus only 5 percent greater than the <u>ab initio</u> PA result.

Second Test

A computational run was then performed on the interaction between BF_3 and NH_3 . NH_3 is a molecule that interacts strongly with surface Lewis acid sites and is widely used as a poison to block the effect of these sites. Optimized geometries are shown in figure 3 and total energy values are listed in table III. Henceforth, only select distances and angles will be reported.

The B-N distance was 1.782 Å, and the net atomic charges on the BF₃ and NH₃ groups were ± 0.326 respectively. The calculated dihedral angle of the BF₃ group, -130.7° , showed this molecule was distorted from its normal planar trigonal geometry. The NH₃ bond angles increased form 109.1° to 111.4° and the dihedral angle increased from

 119.0° to 125.0° . The total energy of the BF₃—NH₃ system was 52.2 kJ/mol lower than the added total energy of the isolated molecules. The HOMO orbital plot for this system is shown in figure 4. One can notice a withdrawal of the nitrogen 2p orbital toward the boron atom.

These results indicated that an adduct was formed between BF₃ and NH₃ which is what one would expect experimentally.

Perfluoro ether Results

(1) BF₃ Interactions

The first two calculations involved the interaction between BF_3 and the O atom of a dimethyl ether molecule and the perfluorodimethyl ether molecule. The optimized structure and total energy for the BF_3 — $(CH_3)_2O$ system are shown in figure 5 and table III. Vibrational analysis of this system revealed no negative frequencies. The distance between the boron atom and the ether oxygen atom was 1.905 Å. The net atomic charge on the BF_3 group was -0.119, and the charge on the dimethyl ether was +0.119. The geometry of the BF_3 group was not planar trigonal as noted its dihedral angle of 159.4°. The total energy of the BF_3 — $(CH_3)_2O$ system was 30.4 kJ/mol lower than the added total energy of the isolated molecules. The HOMO orbital plot for this system showed withdrawal of the oxygen 2p orbital toward the boron atom.

The computed vibrational infrared frequencies, for the BF_3 — $(CH_3)_2O$ adduct, can be compared (see table IV) with the experimental values determined by Hunt and Ault (ref. 11). Very good agreement was observed with the two experimental frequencies assigned to the antisymmetric B—F stretching vibrations (1260 cm⁻¹ and 1221 cm⁻¹) and with the frequency assigned to the B—O stretching vibration (624 cm⁻¹). The computed frequency corresponding to the symmetric B—F stretching vibration (812 cm⁻¹), however, was not observed.

The optimized structure and total energy of the BF_3 —(CF_3) $_2$ O system are shown in figure 6 and table III. Vibrational analysis revealed no negative frequencies. The distance between the boron atom and the ether oxygen atom was 3.594 Å. The net atomic charge on the BF_3 group was zero as well as on the ether group. The geometry of the BF_3 group was planar trigonal as noted by its dihedral angle of 179.8°. The total energy of this system equaled the added total energy of each isolated molecule. The HOMO orbital plot for this system showed no withdrawal of the oxygen 2p orbital towards the boron atom.

A third calculation was performed interacting BF_3 with one of the fluorine atoms (the one with the most negative atomic charge) of the perfluorodimethyl ether molecule. The distance between B and the F atom was 3.557 Å. The net atomic charges on the BF_3 and ether groups were both zero. The geometry of the BF_3 molecule was planar trigonal as noted by the computed angles. The total energy of this system was 0.3 kJ/mol higher than the added total energy of each isolated molecule.

The last computation involved the interaction between AlF_3 and the perfluorodimethyl ether molecule at the oxygen atom of the ether. The distance between the O and Al atoms was 4 Å. The net atomic charges on the AlF_3 and ether groups were both zero. The geometry of the AlF_3 molecule was planar trigonal, and the total energy of this system was 1 kJ/mol higher than the added total energy of each isolated molecule.

These results support the existence of a Lewis acid-base interaction between the BF_3 and dimethyl ether molecules, but no Lewis acid-base interaction between the BF_3 and perfluorodimethyl ether molecules. These computational results are in agreement with the experimental observations of Ball and Zehe (ref. 8). In addition, interaction between the perfluorodimethyl ether and AlF_3 , another Lewis acid, was not noted. Although the presence of the metal atoms of a metal oxide or metal fluoride cannot be ruled out as a contributing factor in the high temperature decomposition of PFPEs, this molecular orbital study does suggest that the formation of a classical Lewis acid-base adduct, like that of the BF_3 — NH_3 system, will not occur between PFPE molecules and surface Lewis acid sites.

CONCLUSIONS

Results of this AM1 semi-empirical molecular orbital study showed the formation of a classical Lewis acid-base adduct between dimethyl ether and BF_3 . The formation, however, of a Lewis acid-base adduct between perfluorodimethyl ether and BF_3 (as well as AlF_3) was not observed, thus confirming the experimental work of Ball and Zehe. Therefore, these results indicate that the high temperature catalytic decomposition of PFPEs is not due to surface Lewis acid sites but to other unknown sites. Future work will focus on identifying these sites.

REFERENCES

- 1. Carré, D.J., and Markowitz, J.A., "The Reaction of PFPE Oil with FeF₃, AlF₃, and AlCl₃ at Elevated Temperatures," ASLE Trans., 28(1), pp. 40–46 (1985).
- 2. Kasai, P.H., Tang, W.T., and Wheeler, P.O., "Degradation of PFPE's Catalyzed by Aluminum Oxide," Applied Surface Science, 51, pp. 201–211 (1991).
- 3. Morales, W., "Perfluoropolyalkylether Decomposition on Catalytic Aluminas," ASA TM-106547 (1994).
- Herrera-Fierro, P., Jones, W.R. Jr., and Pepper, S.V., "Interfacial Chemistry of a Perfluoropolyether Lubricant Studied by X-ray Photoelectron Spectroscopy and Temperature Desorption Spectroscopy," J. Vac. Sci. Technol. A 11(2), pp. 354–367, (1993).
- 5. Zehe, M.J., and Faut, O.D., "Acidic Attack of PFPE Lubricant Molecules by Metal Oxide Surfaces," NASA TM-101962 (1993).
- Basu, P., Ballinger, T.H., and Yates, J.T. Jr., "Fluorinated Ether Chemistry on Alumina: A Transmission IR Study of the Adsorption and Thermal Decomposition of (CF₂H)²O on Al₂O₃," Langmuir, 5(2), pp. 502–510 (1989).
- 7. Ng, L.M., Lyth, E., Zeller, M.W., and Boyd, D.L., "Surface Chemistry of Perfluoro Ethers: An Infrared Study of the Thermal Decomposition of (C₂F₅)₂O on Al₂O₃," Langmuir, 11, pp. 127–135 (1995).
- 8. Ball, D.W., and Zehe, M.J., "Matrix Isolation Studies of the Interactions of BF3 with Water and Substituted Diethyl Ethers," NASA TM-106422, (1993).
- 9. Fontaine, M., Delhalle, J., Defranceshi, M., Lecayon, G., and Boissel, J., "Preliminary Theoretical Study of Perfluorodimethyl Ether and Its Protonated Form," International Journal of Quantum Chemistry, 46, pp. 171–181, (1993).
- 10. Bureau, C., Defranceschi, M., Dehalle, J., and Bourin, J., "Dissociative Chemisorption of Perfluoropolyethers Above Reactive Sites on Nickel Surfaces: A Theoretical Contribution," J. of Molecular Catalysis, 85, pp. 183–200, (1993).
- 11. Hunt, R.L., and Ault, B.S., "Matrix Isolation Spectroscopic Study of the Boron Trihalide-Dimethyl Ether 1:1 Adducts," Int. J. of Spectroscopy, 1, pp. 45–61, (1982).

TABLE I.—PERFLUORODIMETHYL ETHER RESULTS

Bond length, Å	HF/3-21G basis	HF/4-31G basis	AM1 this work	Experimental
(C ₁ -O), (C ₂ -O)	1.370	1.369	1.405	1.369
$(C_1-F_1), (C_2-F_4)$	1.322	1.331	1.351	1.327
$(C_1-F_2), (C_2-F_5)$	1.331	1.342	1.353	1.327
$(C_1-F_3), (C_2-F_6)$	1.337	1.347	1.354	1.327
Bond angles, °				
C ₁ -O-C ₂	124.1	125.9	121.3	119.1
F ₁ -C ₁ -O	108.0	107.6	104.8	
Dihedral angle, °				
F ₁ -C ₁ -O-C ₂		162.0	165.4	164

TABLE II.—PROTONATED PERFLUORODIMETHYL ETHER RESULTS

Bond length, Å	HF/3-21G basis	AM1 this work	
C ₁ -O	1.508	1.521	
Н-О	.989	.992	
C ₁ -F ₁	1.305	1.333	
C ₁ -F ₂	1.301	1.331	
C ₂ -F ₄	1.304	1.333	
Bond angles, °			
C ₁ -O-C ₂	123.0	128.5	
F ₁ -C ₁ -O	103.9	101.9	

TABLE III.—TOTAL ENERGY CALCUALTIONS (kJ/mol)

Molecule	Total energy	System	Total energy
BF ₃ NH ₃ ALF ₃ DME PFDE	-147,795.2 -24,002.0 -145,947.0 -63,664.1 -336,994.1	FB ₃ /NH ₃ BF ₃ /DME BF ₃ /PFDE AIF ₃ /PFDE	-171,849.4 -211,489.7 -484,789.4 -482,940.1

DME—Dimethyl ether

PFDE—Perfluorodimethyl ether

TABLE IV.—COMPUTED VIBRATIONAL INFRARED FREQUENCIES FOR THE $BF_3 - (CH_3)_2O \ ADDUCT$

AM1 this work	Experimental	Assignment
1329 cm ⁻¹	1260 cm ⁻¹	ASYM B-F str.
1205	1221	ASYM B-F str.
	812	SYM B-F str.
637	624	B-O str.




Figure 1.—(a) Initial molecular structure for PFDE. (b) Optimized molecular structure for PFDE.





Figure 2.—Optimized molecular structure for protonated PFDE.



Net atomic charge on NH₃: +0.326 Net atomic charge on BF₃: -0.326

Figure 3.—Optimized molecular interaction between ${\rm BF}_3$ and ${\rm NH}_3.$






Figure 4.—HOMO orbital plot for BF₃–NH₃ interaction.






Net atomic charge on DME: +0.119Net atomic charge on BF₃: -0.119

Figure 5.—Optimized molecular interaction between DME and BF_3 .





Net atomic charge on PFDE: 0 Net atomic charge on BF₃: 0

Figure 6.—Optimized molecular interaction between PFDE and BF_3 .

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED	
	March 1996	Te	chnical Memorandum
4. TITLE AND SUBTITLE	•	·	5. FUNDING NUMBERS
A Molecular Orbital Study on the Dimethyl Ether and Perfluorodic		and the Compounds	
6. AUTHOR(S)			WU-505-63-5A
Wilfredo Morales			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER
National Aeronautics and Space	Administration		
Lewis Research Center			E-10128
Cleveland, Ohio 44135-3191			2 10120
6.6 (S.M.), 6.116 (17)			
9. SPONSORING/MONITORING AGENCY	NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING
			AGENCY REPORT NUMBER
National Aeronautics and Space	Administration		
Washington, D.C. 20546-0001			NASA TM-107174
11. SUPPLEMENTARY NOTES			
II. SOFFLEWENTANT NOTES			
Responsible person, Wilfredo M	Iorales, organization code 51	140, (216) 433–6052.	
12a. DISTRIBUTION/AVAILABILITY STAT	EMENT		12b. DISTRIBUTION CODE
Unclassified - Unlimited			
Subject Categories 27 and 31			
	NIAGAG 4 C A I I C	.: (201) 621 0200	
This publication is available from the 13. ABSTRACT (Maximum 200 words)	NASA Center for Aerospace Info	ormation, (301) 621–0390.	
ether and perfluorodimethyl ether molecule and dimethyl ether but	er. The results indicated the ano interaction with the perfug the interaction between A	existence of a Lewis ac luorodimethyl ether. In	s acid) and the compounds dimethyl id-base interaction between the BF ₃ addition, the results of a second) and perfluorodimethyl ether, also
14. SUBJECT TERMS			15. NUMBER OF PAGES
			9
Quantum chemistry; Lewis acids; Perfluoroethers			
	s; Perfluoroethers		16. PRICE CODE
			A02
	s; Perfluoroethers BECURITY CLASSIFICATION DF THIS PAGE	19. SECURITY CLASSIFICA OF ABSTRACT	A02